



## THYROID



Biological Conflict    Conflict-Active Phase    Healing Phase

**DEVELOPMENT AND FUNCTION OF THE THYROID GLAND:** The thyroid gland is situated at the front of the lower neck below the **larynx** with a right and left lobe on each side of the **trachea**. Originally, the thyroid gland was located in the **oropharynx** from where it descended to its final position, taking a path through the **tongue** and the neck. This connection is known as the **thyroglossal duct**. The primary function of the thyroid is the production of thyroxine (**secretory quality**), a hormone that regulates the rate in which nutrients are converted into energy. Initially, the thyroid was an exocrine gland excreting hormones into the ingoing and outgoing section of the **intestine** to facilitate the ingestion of food and the elimination of feces. After the gullet had broke open, the thyroid became an endocrine gland releasing thyroxine directly into the bloodstream. The thyroid gland consists of **intestinal cylinder epithelium**, originates from the **endoderm** and is therefore controlled from the brainstem.



**BRAIN LEVEL:** In the **brainstem**, the thyroid gland has two control centers that are orderly positioned within the **ring form** of the brain relays that control the organs of the **alimentary canal**.


The right half of the thyroid gland is controlled from the right side of the brainstem; the left half is controlled from the left brainstem hemisphere. There is no cross-over correlation from the brain to the organ.

**NOTE:** The mouth and pharynx, tear glands, Eustachian tubes, thyroid gland, parathyroid glands, pituitary gland, pineal gland, and choroid plexus share the same brain relays.

**BIOLOGICAL CONFLICT:** Consistent with its role in digestion, the **biological conflict** linked to the thyroid gland is a **"morsel conflict"**.


In line with evolutionary reasoning, **morsel conflicts** are the primary conflict theme associated with **brainstem-controlled organs** deriving from the **endoderm**.

### RIGHT HALF OF THE THYROID GLAND



Equivalent to the **right half of the mouth and pharynx**, the conflict linked to the **right lobe of the thyroid** relates to an **"ingoing morsel"** and to **"not being fast enough to catch a morsel"**. Such a "morsel" concerns, for example, a job, a position, a promotion, a contract, a business, or a purchase one strongly desires but is too slow to "grab". The expected "morsel" could also relate to a person one is too slow to "catch" or "get a hold of".

## LEFT HALF OF THE THYROID GLAND



Equivalent to the **left half of the mouth and pharynx**, the conflict linked to the **left lobe of the thyroid** relates to an **“outgoing morsel”** and to **“not being fast enough to eliminate a morsel”** (originally, the feces morsel). This could be a term paper, any kind of goods, foul stocks or a person (tenant, employee, business partner) one was too slow to “get rid of”. An apology or a proposal that was expressed too late can also evoke this type of “morsel” conflict.

People who are driven to “get things done”, who have professions and activities that involve competition (business managers, sales agents, vendors, athletes and sports competitors), who are under deadline pressure (journalists, manufacturers) or constant pressure to “keep up” (working two jobs, working mothers) are more susceptible to experience the conflict. Children and adolescents suffer thyroid-conflicts when they are pushed by a parent, teacher, or coach (“You are too slow!”).

**CONFLICT-ACTIVE PHASE:** Starting with the **DHS**, during the **conflict-active phase**, thyroid gland cells proliferate proportionally to the intensity of the conflict. The **biological purpose of the cell increase** is to improve the production of thyroxine so that the individual becomes faster to catch the desired morsel (right half of the thyroid) or to get rid of an undesired morsel (left half of the thyroid). This causes an **overactive thyroid** or **hyperthyroidism**. Because of the body’s enhanced metabolism, persons with an overactive thyroid are often overexcited, nervous, irritable, and have trouble sleeping.

With persistent conflict activity the growth (**secretory type**) created by the continuing cell augmentation forms a **hard struma**, or **goiter** (compare with **euthyroid struma** related to the **thyroid ducts**). The enlargement of the thyroid could cause breathing difficulties due to the pressure on the **trachea**. A large swelling with profuse cell proliferation might be diagnosed as a **thyroid cancer**.



It is a wide-spread belief that hyperthyroidism is caused by iodine deficiency. This theory, however, cannot explain why, for example, a goiter develops in the right or left thyroid lobe (see picture), or in both.




This CT scan highlights the area in the brainstem from where the left thyroid gland is controlled (**view the GNM diagram**). The **sharp ring configuration** of the **Hamer Focus** indicates conflict activity, hence, an overactive thyroid.

**HEALING PHASE:** Following the **conflict resolution (CL)**, **fungi or mycobacteria** such as TB bacteria remove the cells that are no longer needed. **Healing symptoms** are **pain** due to the swelling, **difficulties breathing and swallowing**, and **night sweats**. If the healing process is accompanied by an inflammation, this causes **thyroiditis**.

With the completion of the healing phase the thyroxine level returns to normal. However, with a **hanging healing**, that is, when healing is continually interrupted by **conflict relapses**, the prolonged decomposing process results in a loss of thyroid gland tissue causing a chronic **underactive thyroid**, or **hypothyroidism**, also termed **Hashimoto’s disease**. **Symptoms** are **fatigue and low energy**, since the insufficient production of thyroxine slows down the body’s metabolism. In this case, supplementing thyroxine is advisable.

**NOTE: Hypothyroidism is always preceded by hyperthyroidism!**

**If the required microbes are not available upon the resolution of the conflict**, because they were destroyed through an overuse of **antibiotics**, the additional cells in the thyroid gland cannot be broken down. Consequently, the growth or goiter stays **maintaining the overproduction of thyroxine** with **lasting hyperthyroidism**, even though the conflict has been resolved (see also **parathyroid glands, pancreas gland, adrenal gland, prostate gland**). To normalize the thyroxine production, surgery might have to be considered.



Biological Conflict

Conflict-Active Phase

Healing Phase

**DEVELOPMENT AND FUNCTION OF THE PARATHYROID GLANDS:** The parathyroid glands are two pairs of small glands located on the back side of the **thyroid gland**. Their main function is to secrete a hormone (PTH-parathyroid hormone) that helps maintain the proper level of calcium (**secretory quality**), a mineral essential for muscle contraction. Like the **thyroid gland**, the parathyroid glands were originally exocrine glands that excreted into the **intestine**. Today, they are endocrine glands that release their hormones directly into the bloodstream. The parathyroid glands consist of **intestinal cylinder epithelium**, originate from the **endoderm** and are therefore controlled from the brainstem.



**BRAIN LEVEL:** In the **brainstem**, the parathyroid glands have two control centers that are orderly positioned within the **ring form** of the brain relays that control the organs of the **alimentary canal**.

The right parathyroid gland is controlled from the right side of the brainstem; the left parathyroid gland is controlled from the left brainstem hemisphere. There is no cross-over correlation from the brain to the organ.

**NOTE:** The mouth and pharynx, tear glands, Eustachian tubes, thyroid gland, parathyroid glands, pituitary gland, pineal gland, and choroid plexus share the same brain relays.

**BIOLOGICAL CONFLICT:** According to the function of the parathyroid glands, the corresponding **biological conflict** is a **"morsel conflict"**.

In line with evolutionary reasoning, **morsel conflicts** are the primary conflict theme associated with **brainstem-controlled organs** deriving from the **endoderm**.

**RIGHT PARATHYROID GLANDS:** Equivalent to the **right half of the mouth and pharynx**, the conflict linked to the right parathyroid glands relates to an **"ingoing morsel"** and to **"not being able to catch a morsel" because of a low calcium level limiting the muscle contraction required to ingest a food morsel**.

**LEFT PARATHYROID GLANDS:** Equivalent to the **left half of the mouth and pharynx**, the conflict linked to the left parathyroid glands relate to an **"outgoing morsel"** and to **"not being able to eliminate a morsel" because of a low calcium level limiting the muscle contraction required to eliminate a morsel**.


**CONFLICT-ACTIVE PHASE:** Starting with the **DHS**, during the **conflict-active phase** cells in the parathyroid glands proliferate causing an **overproduction of PTH** or **hyperparathyroidism** with the **biological purpose** to supply the organism with more calcium to improve the muscular contraction so that the morsel can be better absorbed (right glands) or eliminated (left glands). Consequently, the calcium level in the blood increases causing **hypercalcemia** (compare with **hypercalcemia** related to the **bones**). In conventional medicine, a large growth in the parathyroid glands might be diagnosed as a **parathyroid cancer**.

**NOTE:** The PTH-parathyroid hormone draws the required calcium from the bones. However, this does not cause **osteoporosis**, since PTH ensures at the same time that excess calcium is not excreted through urination but carried back to the organism.

**HEALING PHASE:** Following the **conflict resolution (CL)**, **fungi or mycobacteria** such as TB bacteria remove the cells that are no longer needed. This process is accompanied by **night sweats**. With the completion of the healing phase the PTH level returns to normal. However, with a **hanging healing**, when healing is continually interrupted by **conflict relapses**, the prolonged bacterial activity leads to a loss of parathyroid gland tissue causing chronic **hypoparathyroidism** with constant low calcium levels. In this case supplementation is advisable.


**NOTE:** Hypoparathyroidism is always preceded by hyperparathyroidism!

If the required microbes are not available upon the resolution of the conflict, because they were destroyed through an overuse of antibiotics, the additional cells cannot be broken down causing **lasting hyperparathyroidism** (see also thyroid gland, pancreas gland, adrenal gland, prostate gland). To normalize the thyroxine production, surgery might have to be considered.



**DEVELOPMENT AND FUNCTION OF THE THYROID DUCTS:** The thyroid ducts branch throughout the thyroid gland in a tree-like structure. The original function of the thyroid ducts was to carry hormones produced in the thyroid into the ingoing and outgoing section of the intestine to aid the metabolism of food and the disposal of feces. After the rupture of the gullet, the thyroid ducts closed and the thyroid became an endocrine gland. Today, the thyroid ducts deliver thyroxine directly into the bloodstream. The lining of the thyroid ducts consists of squamous epithelium, originates from the ectoderm and is therefore controlled from the cerebral cortex.


**NOTE:** The thyroid ducts are descendants of the pharyngeal arches (see also coronary arteries, coronary veins, aorta, carotid arteries, and subclavian arteries that derive from the pharyngeal arch arteries. In the embryo, the pharyngeal arches, or branchial arches (Greek branchial = gill), give rise to structures of the head and neck (see also pharyngeal ducts).



**BRAIN LEVEL:** The epithelial lining of the thyroid ducts is controlled from the pre-motor sensory cortex (part of the cerebral cortex). The left thyroid ducts are controlled from the right side of the cerebral cortex; the right thyroid ducts are controlled from the left cortical hemisphere (frontal). Hence, there is a cross-over correlation from the brain to the organ.

**NOTE:** The thyroid ducts and pharyngeal ducts share the same brain relays. The DHS affects either one of the tissues or both, depending on the intensity of the conflict.

**BIOLOGICAL CONFLICT:** The biological conflict linked to the thyroid ducts is a female powerless conflict or male frontal-fear conflict, depending on a person's gender, laterality, and hormone status. A powerless conflict is experienced as feeling helpless ("there is nothing I can do about this", "my hands are tied") or of not being in control of a situation. Generally speaking, the conflict relates to any kind of imposition, external control or decision made over one's head.



The Biological Special Program of the thyroid ducts follows the **GULLET MUCOSA PATTERN** with hypersensitivity during the conflict-active phase and the Epileptoid Crisis and hyposensitivity in the healing phase.

**CONFLICT-ACTIVE PHASE:** ulceration in the lining of the thyroid ducts proportional to the degree and duration of conflict activity. The biological purpose of the cell loss is to widen the ducts to supply the organism with more thyroxine; this provides the individual with more energy to resolve the conflict. **Symptoms:** mild to severe pain, depending on the intensity of the conflict. Since the lumen of the thyroid ducts enlarges, the thyroxine level rises slightly during the conflict-active phase. This, however, must not be confused with hyperthyroidism because the thyroxine production in the thyroid gland is unaffected.

**HEALING PHASE:** During the first part of the **healing phase (PCL-A)** the tissue loss is replenished through **cell proliferation** with **swelling** due to the **edema** (fluid accumulation) in the healing area. Since the thyroid ducts have no external opening, a cyst forms as a result of the back-up of fluid in the duct. Such a thyroid cyst is called a **euthyroid struma**, or **goiter** (compare with **goiter** related to the **thyroid gland**). Since there are no thyroxine-producing cells involved, the thyroxine level stays within the normal range. **Thyroid cysts are located towards the middle (median)** on the right or left side of the neck (compare with cysts in the **pharyngeal ducts** located laterally). If there are no conflict relapses, the swelling recedes in the course of the healing process. However, with a **hanging** healing the cyst stays until healing is complete.




**Thyroglossal cysts** develop in the **thyroglossal duct** that connects the **thyroid gland** with the base of the tongue.



This brain CT presents a **Hamer Focus** on the right side of the cerebral cortex, precisely, in the area from where the left thyroid ducts and thyroglossal duct are controlled (**view the GNM diagram**). The small fluid accumulation, showing as dark, indicates the beginning of PCL-A.



A **thyroid fistula** is an external opening of a **thyroid duct** caused by the rupture of a **thyroid cyst** (euthyroid struma) with fluids emptying outwards. A thyroid cyst can break, for example, when **large amounts of water are retained** in the cyst due to the **SYNDROME** or as a result of continuous **conflict relapses** that prolong the healing process. Yet, a fistula is only created when the right thyroid ducts are affected since they are located closer to the skin. This explains why a **thyroid fistula always forms on the right side of the neck**.



**In the brain**, the right thyroid ducts, where the fistula occurs, are controlled from the left cortical hemisphere exactly opposite the brain relay of the left **thyroid ducts** and the **para-anal ducts**. Here is why: Originally, before the **gullet broke open**, the thyroid was an exocrine gland that released thyroxine into both sections of the **intestine**. The right thyroid ducts (controlled from the left side of the brain) excreted into the ingoing section (today's **mouth and pharynx, esophagus, stomach and duodenum, small intestine**) to aid the digestion of food; the left thyroid ducts (controlled from the right side of the brain) excreted into the outgoing section (today's **rectum**) to accelerate the disposal of feces. However, when the **gullet ruptured**, parts of the left **thyroid ducts** remained in the rectum. These residues are today's **para-anal ducts** (see **para-anal fistula**). The close vicinity of the brain control centers of the thyroid ducts and para-anal ducts represents the **rupture of the gullet** on the cerebral level.